

OSU Veterinary Hospital 601 Vernon Tharp Street Columbus, OH 43210 Phone: (614) 292-3551 Fax: (614) 292-2053	ECHOCARDIOGRAPHY REPORT - CARDIOLOGY SERVICE THE OHIO STATE UNIVERSITY VETERINARY MEDICAL CENTER
	John Bonagura, DVM, DACVIM Karsten Schober, DVM, PhD, DECVIM Jaylyn Rhinehart, DVM, DACVIM Emily Chapel, DVM Michelle Rohrbaugh, DVM Alicia Byrd, RVT Tammy Muse, RVT

Patient Number: 000 **448316**Patient Name: **Sweeney, JD**Date of study: **05/26/2017**

Diagnosing Cardiologist:

Species:

Breed: **Maine Coon**Age: **3**Birthdate: **06/10/2014**Sex: **Male**Weight (kg): **0.0 kg**

BSA:

Systolic BP:

Clinical Findings

The echocardiogram was performed as a screen for hypertrophic cardiomyopathy (HCM) phenotype.

Auscultation: sinus rhythm; no murmurs or gallop sounds.

Screening Exam for Feline Hypertrophic Cardiomyopathy; details: This examination includes subjective evaluation of long and short axis images from the parasternal (intercostal) right-sided acoustic windows. M-mode examination of the LV is also performed. The examination screens for ventricular hypertrophy using 2D long and short axis image planes as well as the standard M-mode images with the cursor placed dorsally to the posterior papillary muscle. Left atrial size is also assessed subjectively and by long-axis maximal diameter. Doppler studies are only performed if needed to evaluate gallop sounds or any murmurs if present.

Echocardiographic Findings

The echocardiographic examination was conducted from the right side of the thorax. The technical examination was of diagnostic quality and the patient was sufficiently cooperative.

Normal 2D & M-mode Study (Doppler studies not performed):

There were no congenital or acquired structural cardiac lesions observed by 2D echocardiography.

All cardiac chambers and great vessels were within normal size limits.

There were no overt valvular lesions.

Left ventricular ejection fraction (shortening fraction) was normal.

Diagnosis & Recommendations

Normal screening examination; no evidence fo hypertrophic cardiomyopathy or congenital disease - similar to prior exam (JDB)+WNL

<u>2D Measurements</u>		<u>M-Mode</u>		<u>Doppler Measurements</u>
LA Diam	19.8 mm	IVSd	4.4 mm	
LA2D/LVIDd	1.1 (0.8 - 1.1)	LVIDd	18.4 mm	
IVSd-max-Laxis	4.9 mm	LVPWd	5.1 mm	
LVPWd-max-Laxis	5.1 mm	IVSs	7.1 mm	
		LVIDs	10.2 mm	
		LVPWs	8.6 mm	
		EDV(Teich)	10.3 ml	
		ESV(Teich)	2.1 ml	
		EF(Teich)	79.2 % (> 48.0)	
		%FS	44.9 % (> 25.0)	
		SV(Teich)	8.18 ml	

Echocardiogram Reported by: John D. Bonagura, DVM, DACVIM (Cardiology, Internal Medicine) _____